

Histocompatibility Antigens in Patients with Leprosy from Azarbaijan, Iran¹

N. Mohaghehpour, H. Tabatabai, K. Mohammad,
K. Ramanujam, and F. Z. Modabber²

Evidence in man indicates that the clinical picture in leprosy as well as the prognosis for patients with this disease parallels the efficacy of the host's cellular immune response (⁶). In lepromatous leprosy there is a marked deficiency of the cell-mediated immune response (CMI); such impairment is not a general feature of the tuberculoid form. The documented deficiency of CMI in patients with lepromatous leprosy has been suggested to be a consequence of intensive exposure to *Mycobacterium leprae* (⁷); however, the possibility that such a deficiency antedates infection cannot be overlooked. The familial pattern of leprosy as well as the high incidence in certain genetically-related populations have implicated a genetic predisposition for susceptibility to this disease (¹).

Recent evidence concerning the association between susceptibility to certain diseases, major histocompatibility antigens, and immune responses (^{9, 11, 13, 15}) warrants a search for the possible association of HLA antigens with manifestation of leprosy. Thus, in this study the HLA profiles of 88 Iranian patients with leprosy were compared with those of normal individuals from corresponding ethnic groups.

MATERIALS AND METHODS

Patients. Eighty-eight leprosy patients included in this investigation were from

Baba Baghi Leprosarium in Western Azarbaijan, Iran. Patients were characterized by clinical, bacteriological, and histological examination. Sixty-five patients had the lepromatous form of leprosy (LL), 9 cases were borderline (BB), 7 were borderline with lepromatous features (BL), and 7 were indeterminate (I), using the Ridley and Jopling system (¹⁹).

All patients were from Eastern Azarbaijan. Fifty-nine patients were linguistically Turkish (Azeri Turkie) and 29 were Kurdish. Controls were 125 normal individuals from the same regions as the patients, including 101 Turks and 24 Kurds.

Serological methods. HLA antigens were identified by the standard microlymphocytotoxicity test (²²), using typing trays obtained from Terasaki. Upon receipt, reactivity of the reagents was checked using HLA sera. Because of serological uncertainties, antigens A₉, A₁₀, AW₁₉ and BW₁₆ were not subdivided.

Data analysis. Statistical tests were performed using the Mantel-Haenszel procedure (¹²) to provide a one degree of freedom χ^2 for the differences in antigen frequency between patients and controls across the different ethnic groups. Adjustment to the nominal significance levels was made to account for the number of different comparisons being made. The ordinary chi square test was used for other comparisons. Relative risk (x) was calculated for significant associations according to the method described by Woolf (²⁵).

RESULTS

In the two ethnic groups, Kurds and Turks, comparison of the frequencies of HLA antigens using the Mantel and Haenszel method did not reveal any differences between the patients with lepromatous leprosy or the total patient populations and the controls (The Table).

¹ Received for publication on 3 July 1979.

² N. Mohaghehpour, Ph.D., Associate Professor, Department of Pathobiology, School of Public Health, H. Tabatabai, M.Sc., Research Assistant, Department of Pathobiology, School of Public Health, Teheran University, Teheran, Iran; K. Mohammad, Ph.D., Assistant Professor, Department of Epidemiology and Biostatistics, Teheran University; K. Ramanujam, M.D., Professor, Schieffelin Leprosy Research and Training Centre, Karagiri, North Arcot, Tamil Nadu, South India; F. Z. Modabber, Ph.D., Associate Professor, Department of Pathobiology and Institute of Biophysics and Biochemistry, Teheran University.

TABLE. Comparison of HLA antigen frequencies (%) among leprosy patients and healthy controls.

Antigen	Kurds			Turks		
	All patients (29) ^b	LL. ^a patients (20)	Controls (24)	All patients (59)	LL patients (45)	Controls (101)
A ₁	10	5	29	15	16	17
A ₂	21	20	16	25	20	26
A ₃	24	30	25	20	24	20
A ₉	38	40	38	20	20	26
A ₁₀	14	10	29	10	11	7
A ₁₁	21	20	4	31	31	22
A ₂₈	0	0	8	7	9	12
AW ₁₉	31	35	25	34	38	29
B ₅	28	25	38	39	38	34
B ₇	3	0	4	5	4	7
B ₈	10	15	4	10	11	6
B ₁₂	21	15	17	12	13	14
B ₁₃	3	5	0	10	7	6
B ₁₄	3	5	4	3	4	4
B ₁₈	14	15	13	8	11	7
B ₂₇	3	5	0	2	2	5
BW ₁₅	7	10	21	2	2	10
BW ₁₆	14	15	33	12	16	5
BW ₁₇	7	5	0	5	6	4
BW ₂₁	7	5	8	2	2	8
BW ₂₂	17	15	8	14	11	10
BW ₃₅	31	40	8	36	36	35
BW ₃₇	3	5	4	0	2	1
BW ₄₀	3	0	8	12	13	8

^a LL, lepromatous leprosy.

^b Number of individuals tested.

Looking at the different groups individually, Kurdish speaking patients with the lepromatous form of the disease (LL patients) showed an increase in HLA-BW₃₅ frequency (40%) as compared to the normal individuals (8%) ($p < 0.02$, $x = 7.3$) (The Table). The frequency of this antigen in the total patient population was also higher than that observed among the healthy Kurds ($p < 0.05$, $x = 5.0$). Moreover, HLA-A₁ showed a diminished frequency among the LL patients as compared to the controls ($p < 0.05$, $x = 0.13$). None of these comparisons, however, reached statistical significance when p values were corrected for the number of specificities (twenty-four) tested.

Comparison of the distribution of HLA antigens among the Turks revealed notable deviation in the percentage of HLA-BW₁₅, with 1/59 in patients and 10/101 (10%) in the controls ($p < 0.05$, $x = 0.16$). However, when corrected for the number of antigens

tested, again, this difference was not statistically significant.

DISCUSSION

In view of the complexity of the genetic composition of the present day population in Iran, the HLA profiles of 88 leprosy patients from Azarbaijan Province were compared to those of 125 healthy, ethnically matched individuals. Among the Kurds, frequencies of HLA-A₁ and HLA-BW₃₅ in patients with lepromatous leprosy were found to be different, with statistical significance at the 5% and 2% levels respectively from the percentage of these antigens in the healthy controls, without correction for the number of antigens tested. Differences in the distribution of HLA-BW₃₅ remained significant when the total patient population was compared with the control group (uncorrected $p < 0.05$). The only association observed in the Turkish population was between leprosy and HLA-BW₁₅ (uncorrected

$p < 0.05$). Comparison of the distribution of HLA antigens across the two ethnic groups, on the other hand, revealed only equivocal differences between the LL patients and the controls. Similarly frequencies of HLA antigens did not show a significant difference when the total patient population was compared with healthy individuals.

An earlier report from Iran has revealed a higher, but statistically insignificant, percentage of HLA-A₁₁ antigen in patients with leprosy (¹⁴). Among the patients examined in the present investigation, the frequency of this antigen was only marginally elevated. It should be noted though that the population tested by Massoud, *et al.* (¹⁴) is not from a distinct ethnic group.

Including the present report, 13 groups have studied the distribution of HLA antigens among patients with leprosy (3, 5, 8, 10, 14, 16, 17, 18, 20, 21, 23, 24); significant associations were shown in six studies (5, 8, 10, 16, 23, 24). Such significant findings are strongly suggestive of a possible association between HLA antigens and susceptibility to leprosy, especially with a particular form of the disease, and cannot be regarded as a chance event. Moreover, family studies by de Vries, *et al.* (⁴) have clearly indicated that both susceptibility to and type of leprosy are controlled by at least two HLA-linked genes. In addition, twin studies provide strong evidence for some type of genetic influence in the predisposition towards leprosy (²). In view of these findings, studies of additional cases and controls and particularly haplotype frequencies seems to be justified.

SUMMARY

The distribution of 24 histocompatibility antigens in 88 Azerbaijani patients with leprosy was determined and compared with those of 125 normal, ethnically matched individuals. HLA-BW₃₅ was increased in frequency among the Kurdish patients as compared to the controls; HLA-A₁₁, however, displayed decreased frequency in patients with the lepromatous form of the disease. Among the Turks, diminished frequency of HLA-BW₁₅ was noted in the total patient population. None of these comparisons, however, reached statistical significance when corrected for the number of antigens

tested. Across the two ethnic groups, differences in the frequencies of HLA antigens between the patients and the controls were only marginal.

RESUMEN

Se estudió la distribución de 24 antígenos de histocompatibilidad en 88 personas Azerbaijani con lepra y en 125 individuos sanos correspondientes al mismo grupo étnico. Comparando los resultados con aquellos encontrados en los controles, se observó que el antígeno HLA-BW₃₅ tuvo una frecuencia más elevada en los pacientes Kurdos mientras que el HLA-A₁₁ tuvo una frecuencia disminuída en los pacientes con la forma lepromatosa de la enfermedad. Entre los Turcos, se observó una frecuencia disminuída del HLA-BW₁₅ en la población total de enfermos. Ninguna de estas comparaciones, sin embargo, alcanzó significación estadística después de haber corregido los datos por el número de antígenos estudiados. Las diferencias en las frecuencias de los antígenos HLA en los pacientes y sus controles entre los dos grupos étnicos (Kurdos y Turcos) fueron tan sólo marginales.

RÉSUMÉ

La distribution de 24 antigènes d'histocompatibilité a été déterminée chez 88 malades de l'Azerbaïdjan et comparée avec celle de 125 individus normaux, assortis sur le plan ethnique. L'antigène HLA-BW₃₅ présentait une fréquence augmentée chez les malades kurdes, comparée aux témoins. Néanmoins, la fréquence de HLA-A₁₁ était diminuée chez les malades atteints de la forme tuberculoïde. Chez les Turcs, dans l'ensemble des malades, on a observé une diminution de HLA-BW₁₅. Aucune de ces comparaisons, toutefois, n'a atteint un seuil de signification statistique, lorsque l'on prenait en compte le nombre d'antigènes étudiés. Les différences notées dans les fréquences d'antigènes HLA chez les malades et chez les témoins, dans les deux groupes ethniques, n'étaient que limites.

Acknowledgments. We are grateful to those who generously donated blood. We thank Mrs. G. Bahrami and Mr. H. Alizadeh for invaluable assistance in recruiting blood donors, Miss M. Kamali for technical assistance, and Miss E. Betsayyad for excellent secretarial help. We also acknowledge the continued cooperation of the physicians and staff of Baba Baghi Leprosarium and thank Dr. S. Sadeghi for classification of the patients.

This investigation received financial support from the World Health Organization. In addition, it was supported in part by funds from the Research Council of the University of Teheran.

REFERENCES

1. BLUMBERG, B. S. Leprosy research through genetics. *Int. J. Lepr.* **33** (1965) 739-743.
2. CHAKRAVARTI, M. R. and VOGEL, F. A. A twin

- study on leprosy. In: *Topics in Human Genetics*, Vol. 1, P. E. Becker, W. Lenz, F. Vogel and G. G. Wendt, eds., Stuttgart: Georg Thieme, 1973, 124 pp.
3. DASGUPTA, A., MEHRA, N. K., GHEI, S. K. and VAIDYA, M. C. Histocompatibility antigens (HL-A) in leprosy. *Tissue Antigens* **5** (1975) 85-87.
 4. DE VRIES, P. R. P., LAI A. FAT, R. F. M., NIJENHUIS, L. E. and VAN ROOD, J. J. HLA-linked genetic control of host response to *Mycobacterium leprae*. *Lancet* **2** (1976) 1328-1330.
 5. ESCOBAR-GUTIERREZ, A., GORODEZKY, C. and SALAZAR-MALLEN, M. Distribution of some of the HL-A system lymphocyte antigens in Mexicans. *Vox Sang.* **25** (1973) 151-155.
 6. GODAL, T., MYRVANG, B., STANFORD, J. L. and SAMUEL, D. R. Recent advances in the immunology of leprosy with special reference to new approaches in immunophylaxis. *Bull. Institut. Pasteur* **72** (1974) 273-310.
 7. GODAL, T. and NEGASSI, K. Subclinical infection in leprosy. *Br. Med. J.* **3** (1973) 557-559.
 8. GREINER, J., SCHLEIERMACHER, D., SMITH, T. L., LENHARD, V. and VOGEL, F. The HLA system and leprosy in Thailand. *Hum. Genet.* **42** (1978) 201-213.
 9. KATZ, S. I., FALCHUK, Z. M., DAHL, M. V., ROGENTINE, G. H. and STROBER, W. A genetic link between dermatitis herpetiformis and gluten-sensitive enteropathy. *J. Clin. Invest.* **51** (1972) 2977-2980.
 10. KREISLER, M., ARNAIZ, A., PEREZ, B., CRUZ, E. F. and BOOTELLO, A. HL-A antigens in leprosy. *Tissue Antigens* **4** (1974) 197-201.
 11. MACKAY, I. R. and MORRIS, P. J. Association of autoimmune active chronic hepatitis with HL-A_{1,8}. *Lancet* **2** (1972) 793-795.
 12. MANTEL, N. and HAENSZEL, W. Statistical aspects of the analysis of data from retrospective studies of disease. *J. Nat. Cancer Inst.* **22** (1959) 719-768.
 13. MARSH, D. G., BIAS, W. B. and HSU, S. H. Association of the HL-A₇ cross-reacting group with a specific reaginic antibody response in allergic man. *Science* **179** (1973) 691-693.
 14. MASSOUD, A., NIKBIN, B., NAZARI, G. R., SYADAT, N. A. and ALA, F. A study of cell-mediated immunity and histocompatibility antigens in leprosy patients in Iran. *Int. J. Lepr.* **46** (1977) 149-153.
 15. MCDEVITT, H. O. and BODMER, W. F. HL-A, immune response genes and disease. *Lancet* **1** (1974) 1269-1275.
 16. NAKAJIMA, S., KOBAYASHI, S., NOHARA, M. and SATO, S. HLA antigen and susceptibility to leprosy. *Int. J. Lepr.* **45** (1977) 273-277.
 17. REA, T. H., LEVAN, N. E. and TERASAKI, P. I. Histocompatibility antigens in patients with leprosy. *J. Infect. Dis.* **134** (1976) 615-618.
 18. REIS, A. P., MAIA, F., REIS, V. F., ANDRADE, I. M. and CAMPOS, A. A. HLA antigens in leprosy. *Lancet* **2** (1974) 1384.
 19. RIDLEY, D. S. and JOPLING, W. H. Classification of leprosy according to immunity. A five group system. *Int. J. Lepr.* **34** (1966) 255-260.
 20. SMITH, G. S., WALFORD, R. L., SHEPARD, C. C., PAYNE, R. and PROCHAZKA, J. Histocompatibility antigens in leprosy. *Vox Sang.* **28** (1975) 42-49.
 21. TAKATA, H., SADA, M., OZAWA, S. and SEKIGUCHI, S. HLA and mycobacterial infection: Increased frequency of B₈ in Japanese leprosy. *Tissue Antigens* **11** (1978) 61-64.
 22. TERASAKI, P. and MCCLELLAND, J. D. Microdroplet assay of human serum cytotoxins. *Nature* **204** (1964) 998-1000.
 23. THORSBY, E., GODAL, T. and MYRVANG, B. HL-A antigens and susceptibility to disease. II. Leprosy. *Tissue Antigens* **3** (1973) 373-377.
 24. YOUNGCHAIYUD, U., CHANDANAYINGYONG, D. and VIBHATAVANAJA, T. The incidence of HLA antigens in leprosy. *Vox Sang.* **32** (1977) 342-345.
 25. WOOLF, B. On estimating the relation between blood group and disease. *Ann. Hum. Genet.* **19** (1955) 251-253.