



Mechanism of Action of the Folate Blocker Diaminodiphenylsulfone (dapsons, DDS) Studied in *E. coli* Cell-Free Enzyme Extracts in Comparison to Sulfonamides (SA)¹

J. K. Seydel, M. Richter, and E. Wempe²

The exceptionally high antibacterial activity of DDS against *M. leprae* (12, 18) leads to the question of the underlying molecular mechanism. The special situation in leprosy, i.e., the lack of an *in vitro* system to cultivate *M. leprae*, forced us to study the mechanism of action and structure-activity relations of sulfones in model strains in order to search for the reasons for their extremely high activity against *M. leprae* (0.0016 $\mu\text{mol/l}$ in the mouse foot pad (11, 12, 18, 28)). In studies with other bacterial strains, it has been found that sulfonamides (SA) (I) and sulfones (DDS) (II) act as "folate blockers."



Recognition of the importance of p-aminobenzoic acid (PABA) in folic acid metabolism and the ability of certain folate derivatives and/or products to reverse the effect of SA or sulfones has led to a general theory of SA and DDS action. In principle, this theory states that SA and DDS are competitive inhibitors of the enzymatic incorporation of PABA in folic acid synthesis (20). The sequential pathway of folic acid synthesis has been evaluated by Jaenicke and Chan (9), Brown (4, 5), Shiota, *et al.* (29), and Ortiz and Hotchkiss (16) and is given in Fig. 1. It has also been shown that bacterial- (16, 31), plant- (8, 15), and plasmodial- (6)

cell-free folate synthesizing extracts are inhibited by SA.

MATERIALS AND METHODS

Determination of DDS and sulfonamide inhibitory activities i_{50} and k_{50} in a cell-free system. The ratio of the amount of folate produced in the reaction mixtures (reactants and concentrations as described by Miller, *et al.* (14)) containing DDS or sulfonamides respectively to that produced in controls was used as fractional activity a . The concentration of inhibitors causing a 50% inhibition of folate synthesis was obtained by plotting $i = 1 - a$ vs. the logarithm of the inhibitor concentration (DDS or SA). In kinetic experiments the rate of folate synthesis was evaluated graphically from the initially linear portions of plots of the amount of folate synthesized vs. time. The concentration of DDS or SA causing a 50% inhibition in the rate of folate synthesis was evaluated from plots of $1/(k_0 - k_i)$ vs. the reciprocal of the inhibitor concentration where k_0 and k_i are the rate constants for folate synthesis in the absence and presence of inhibitor. For a detailed description, see Miller, *et al.* (14).

Assay of folate growth equivalents. The methods have already been described (14).

Minimum inhibitory concentrations (MIC). The methods for MIC determination have been previously described (10, 23, 24).

QSAR analysis. Statistical analysis was performed with a Wang computer 700B or 2000 VP. For all regression equations the number of data points or number of compounds used is n ; the correlation coefficient r , the standard deviation s , and the F-test are given.

¹ Received for publication on 24 September 1979; accepted for publication on 10 December 1979.

² J. K. Seydel, Dr. rer. nat., Dipl. Chem., Professor and Head; M. Richter, Chemical Assistant; E. Wempe, Chemical Assistant; Biochemistry Department, Borstel Research Institute, D-2061 Borstel, Germany.

Pteridine derivatives. The pteridine analog of DDS 6-(N-4,4'-diaminodiphenylsulfonyl) methyl-pteridine has been synthesized in accordance with the description already published for the SA analog N'-d-(5-methylisoxazolyl)-N⁴-(6-pteridinylmethyl) sulfanilamide (3).

nmr (DMSO, 90 MHz) δ 8.72 (S 1H)
AA'BB' centered at 6.65 and 7.53 (J = 8Hz, 8) 4.52 (S 2H)
uv (0.1 N NaOH) 365 nm (ϵ = 6.84), 300 nm (ϵ = 19.05), 225 nm (ϵ = 26.38)

Enzymatic reaction mixtures for the determination of reaction rates for DDS and SA incorporation into a dihydrofolic acid analog. The reactants and concentrations were as described previously (2) with the exception that PABA derivatives were replaced by DDS or SA.

Bacterial growth kinetics. *E. coli* (mutaflo) maintained on agar slants was used as the test organism. The culture broth was dextrose-salts-casamino acids (vitamin-free), which has been described by Anton (1). Trimethoprim (TMP) was supplied by Deutsche Wellcome GmbH, Grossburgwedel; PABA and DDS were from commercial sources.

Growth. A broth culture was inoculated from an agar culture and allowed to grow for 12–16 hr. All cultures were grown at 37°C. A dilution of this culture—the preliminary culture—was allowed to grow into the logarithmic phase. When a concentration of about 10⁷–10⁸ organisms/ml was attained, this culture was used to prepare experimental probes (10⁴ organisms/ml).

Total count (Coulter Counter). Samples of the experimental cultures were diluted with particle free saline (0.85%)-formaldehyde (0.2%) solutions so that a count of 1000–20,000 organisms was obtained. Diluted samples were counted with a Coulter Counter model "ZB" equipped with a 30 μ m orifice. Counts per 50 μ l were obtained. Instrument settings were: 1/aperture current 1; 1/amplification 1/2; matching switch 40 K; gain 10; lower threshold 7 and upper threshold maximum (26, 27). Time of drug addition was as indicated.

RESULTS AND DISCUSSION

Determination of cell-free activity (i₅₀) of SA and DDS. Brown (3) has determined inhibitory activities of several SA in such a cell-free folate synthesizing system. He re-





FIG. 1. Reaction scheme of 7,8-dihydropteroic acid (III) biosynthesis.

ported that these activities in general are proportional to the antibacterial activities (minimal inhibitory concentrations or MIC) of these compounds. The relatively small number and the heterogeneity of the studied compounds, however, have limited utility for detailed quantitative structure-activity relationships (QSAR). Similarly, the data have limited utility in assessing the role of permeability as it relates to antibacterial activity.

To gain more information about structural requirements and the physicochemical forces which are directly related to the enzyme-SA or enzyme-DDS interaction and to find an explanation for the observed optimum in whole cell activity (MIC), the inhibitory activity of DDS and of a closely homologous series of SA¹⁴ have been studied in our laboratories. Results of some typical (single point) experiments conducted to determine the inhibitory activities of several N¹-phenyl (III) and N¹-pyridyl (IV) sulfanilamides and DDS (II) are given in Figs. 2 and 3.



They show a plot of fractional inhibition of folate synthesis against the inhibitory concentration used, i.e., dose response curves. The amount of folate synthesized was determined after a fixed time interval (3 hr). From such plots the SA or DDS concentration was determined which caused 50% inhibition of folate synthesis (i₅₀). The tested SA (\approx 50 in total) show a wide range in physicochemical properties (e.g., lipophilicity π , log P), electron distribution (pKa), and steric effects (E_s) and exhibit an ap-




FIG. 4. Relationship of cell-free inhibitory activities (i_{50}) to whole cell inhibitory activities (MIC, *E. coli*). For discussion of deviating compounds 43–45 and 47 see text. (14) Numbers designate different compounds.

a special “steric” effect in case of the *o*-substituted SA, a dummy parameter, *D*, has been introduced into the regression equation (2) having the value 1 for *o*-substituted derivatives and 0 for *m*- and *p*-substituted SA. The following equation was obtained:

$$\log \text{MIC}_{o,m,p} = 0.67\text{pKa} - 0.24\text{D} - 4.74 \quad \begin{matrix} n & r & s & F \\ 18 & 0.95 & 0.14 & 67 \end{matrix} \quad (2)$$

A similar analysis to explain the observed variance in cell-free activities (i_{50}) resulted in the following equation:

$$\log i_{50,o,m,p} = 0.46\text{pKa} - 0.33\text{D} - 2.47 \quad \begin{matrix} n & r & s & F \\ 18 & 0.97 & 0.08 & 123 \end{matrix} \quad (3)$$

The dependencies on structural changes are almost identical in both systems, thus supporting the assumption that the interaction with the enzyme proteins is the rate determining step (associated with the rate constant k_2 in Fig. 5).

Kinetics of inhibition and formation of analogs. The deviation of compounds 43, 44, 45 and 47 in Fig. 4 leads us to an interesting observation. These compounds, which show more potent inhibitory activities in a cell-free system (i_{50}) than in whole cells (MIC) (Fig. 4) but still less than expected according to equation 3, have very low pKa values (≤ 6.5), i.e., they are more than 90% ionized under the conditions of the experiment. For these compounds the kinetics of




FIG. 5. Schematic drawing of possible rate determining steps in antibacterial action.

inhibition was determined in a cell-free system. An example is given in Fig. 6 for DDS. The observed k_{50} values (concentration




FIG. 6. Determination of inhibitory activity of DDS by the kinetic method.




FIG. 7. Kinetics of inhibition of folate synthesis by DDS ($25 \mu\text{M}$) with (\blacktriangle) and without (\circ) preincubation. In the case of preincubation $25 \mu\text{M}$ DDS were preincubated for 3 hr with 7,8-dihydro-6-hydroxymethylpterin ($32 \mu\text{M}$), and PABA ($20 \mu\text{M}$) was subsequently added.

necessary to reduce the rate of dihydrofolic acid synthesis to one half) are in agreement with the observed i_{50} values (concentration needed to reduce the amount of folate produced after 3 hr to one half). For the very active SA, however, the k_{50} was much smaller than the i_{50} for the first 2–3 hr interval. Later, the rate of folate synthesis returned to the rate of the control probe. This behavior indicates a "use up" of inhibitor in a reaction with the substrate dihydropteridine alcohol (¹⁴). In the case of SA this has been demonstrated, and the dihydroptericoic acid analog has been isolated and identified from cell-free reaction systems and from whole cells (^{3, 25}).

Studies performed in our laboratories revealed that in the case of DDS, an analog of dihydroptericoic acid is formed. If in a cell-free folate synthesizing enzyme system





FIG. 8. NMR spectra of DDS and its dihydropteridinyl analog (solvent DMSO).

the substrate dihydropteridine alcohol pyrophosphate is incubated together with DDS in the absence of competing PABA, no production of dihydroptericoic acid occurs (Fig. 7) when PABA is subsequently added because all precursor dihydropteridine alcohol pyrophosphate has reacted with DDS to form the analog. The dihydroptericoic acid analog—where the PABA moiety has been replaced by DDS—has been synthesized and the structure identified by NMR spectroscopy (Fig. 8). The biochemically synthesized product is identical with the synthetic product.



To examine if the rate determining step in inhibition of folate synthesis by SA and sulfones is the competition for binding sites or the rate of formation of the dihydroptericoic acid analog, associated with the rate constant k_3 in Fig. 5, the kinetics of analog formation was studied in the absence of PABA. The decrease in substrate concen-




FIG. 9. Rate of formation of "folate analogs" by various SA and DDS in absence of PABA (cell-free folate synthesizing enzyme system *E. coli*).

tration (dihydropteridine alcohol) and the formation (increase) of analog has been followed by quantitative thin layer chromatography (²). The results are given in Table 1 and Fig. 9. All SA derivatives studied and DDS show the same rate of analog formation despite the large variance in their inhibitory power (only PABA shows a faster rate for its incorporation into dihydroptericoic acid). Therefore the rate determining step in folate inhibition is not the rate of




FIG. 10. Reversal of DDS inhibited folate synthesis by increasing concentrations of PABA (cell-free folate synthesizing system *E. coli*).

formation of the dihydroptericoic acid analogs but the different affinity of the inhibitory SA and sulfones to the enzyme dihydroptericoic acid synthetase compared to the affinity of the natural metabolite PABA expressed by the Michaelis-Menten constant K_m . The molar concentration of PABA needed to reduce the inhibitory effect of DDS (Fig. 10) and of various SA (at constant concentration of the ionized form) to

TABLE 1.

Rate of H ₂ -ptericoic acid analog formation by various N ¹ -phenylsulfanilamides (SA) in a cell-free system in the absence of PABA at pH 7.71, 37 °C									
$\text{H}_2\text{N}-\text{C}_6\text{H}_4-\text{SO}_2-\text{NH}-\text{C}_6\text{H}_4-\text{R}$	SA-conc. [μM]		MIC [μM]	i ₅₀ [μM]	rate [mm ² min ⁻¹]	pKa	π	D	
	total	ion. fract.							
4-CH ₃	5239	223	21.8	—	6.50	9.25	0.44	0	
H	2840	223	16.0	45.0	7.40	8.97	0	0	
4-Cl	1242	223	13.0	35.0	7.58	8.56	0.83	0	
4-Cl	273	49	13.0	35.0	7.90	8.56	0.83	0	
2-Cl	991	223	2.8	13.5	6.35	8.18	0.37	1	
2-Cl	341	117	2.8	13.5	6.65	8.18	0.37	1	
2-NO ₂	250	205	1.0	4.0	7.90	7.24	0.12	1	
4-NO ₂	250	223	1.0	7.0	7.70	6.97	0.59	0	

PABA	250	250	—	—	14.00	4.67	—	—	
DDS	250	0	16.0	32.0	7.50	- log P 0.9		0	

TABLE 2.

Competitive activity of PABA (K_{PABA}) against various N^1 -phenylsulfanilamides (SA) at constant concentration of ionized SA at pH 7.71, 37°C, in a cell-free system (PABA concentration range 10–50 μ M)								
$H_2N-C_6H_4-SO_2-NH-C_6H_4-R$	SA-conc. [μ M]		MIC	i_{50}	K_{PABA}	pKa	π	D
	total	ion. fract.	[μ M]	[μ M]	[μ M]			
4-OCH ₃	436.7	10	34.5	75.0	59	9.34	-0.12	0
H	185.2	10	16.0	45.0	68	8.97	0	0
4-Cl	80.0	10	13.0	35.0	59	8.56	0.83	0
4-J	38.6	10	11.3	25.0	63	8.17	1.12*	0
4-COCH ₃	16.3	10	2.0	10.5	54	7.52	-0.55*	0
4-NO ₂	12.0	10	1.0	7.0	65	6.97	0.59	0
2-Cl	39.4	10	2.8	13.5	93	8.18	0.37	1
2-NO ₂	13.4	10	1.0	4.0	100	7.24	0.12	1
DDS	50	0	16.0	35.0	16	-	log P 0.9	0

* literature value

one-half of the effect in the absence of PABA is given in Table 2 and Fig. 10 (cell-free system). It is obvious that the ionized form of SA is a more powerful inhibitor of folate synthesis than dapsone. This is in agreement with the other results obtained with *E. coli*, especially the MIC.

So far, the results of mechanism of action studies of sulfones in a cell-free system and in whole cell systems of *E. coli* and also in plasmodia seem to indicate identical inhibitory behavior compared to SA. Also, in a strain of mycobacteria (*M. kansasii*) where DDS shows a remarkable increase in inhibitory power (MIC 0.3–0.5 μ g/ml = 1.2–2.0 μ mol/l), the inhibitory effect can be totally antagonized by PABA (PABA:DDS = 2:1) (17).

Bacterial growth kinetic studies. To get further information which might indicate a deviation in the mechanism of action of DDS in comparison to SA—at least to a certain degree—bacterial growth kinetic studies have been performed (7, 27). In these studies the rate of growth of bacteria is determined during the exponential growth phase in the absence and presence of inhibitors. The number of organisms in a defined volume of culture medium is counted with a Coulter Counter as a function of time and

inhibitor concentration. If the number of bacteria (N) per volume is plotted against time (t) in a semilogarithmic graph, a straight line is obtained:

$$\log N = \log N_0 + \frac{kt}{2.303} \quad (4)$$

From the slopes of these curves the generation rate constant (k) can be calculated. Figure 11 gives an example of the growth of *E. coli* in the absence and presence of various concentrations of DDS. In agreement with the results in the case of SA (27), a lag phase of approximately 5 generation times is observed before onset of inhibition. The lag phase is independent (sampling every 30 min) of the type and concentration of the SA or sulfone used and is due to a pool of dihydrofolate within the bacterial cell. After depletion of this pool, a new steady state growth as a function of inhibitor concentration is obtained. It is interesting to note that a second stronger inhibition phase is observed after a constant time interval if the cultures are treated with DDS. The reason for this is not yet understood. This point will be stressed later.

The relation between concentration and inhibitory effect (decrease in generation rate constant) can be linearized by the fol-




FIG. 11. Typical generation rate curves of *E. coli* (mutafloor) at 37°C in the presence of various concentrations of DDS. The curves, final drug concentrations ($\mu\text{mol/l}$) and apparent steady state rate constants, $k_{\text{app}} \times 10^{-4}$ in sec^{-1} were: (x) control 3.63; (O) 5 DDS, 3.43; (●) 10 DDS, 3.09; (Δ) 15 DDS, 2.8 and for the second inhibition phase: (O) 5 DDS, 3.13; (●) 10 DDS, 2.45; (Δ) 15 DDS, 1.97; (\blacktriangledown) 20 DDS, 1.71; (\odot) 25 DDS, 1.39.

lowing equation (Lineweaver-Burk-Plot):

$$1/(k_0 - k_{\text{obs}}) = \frac{1}{C_{\text{DDS}}} K_A + K_B$$

where k_0 is the rate constant of the control culture, k_{obs} the rate constant in the presence of DDS, C_{DDS} is the DDS concentration, K_A the activity constant obtained from the slope of a curve where $1/(k_0 - k_{\text{obs}})$ is plotted against $1/C$ (see Fig. 12), and $K_B = 1/k_0$ for $C \rightarrow \infty$. The obtained activity constant for DDS was $K_{\text{A}_{\text{DDS}}} = 9.02 \times 10^{-2} \text{ M} \cdot \text{sec} \cdot \text{l}^{-1}$ and for 3-sulfa-5-methylisoxazole $K_{\text{A}_{\text{SMZ}}} = 0.425 \times 10^{-2} \text{ M} \cdot \text{sec} \cdot \text{l}^{-1}$. The ratio of the activity constant $K_{\text{A}_{\text{SMZ}}}$ for these two compounds is the same as the ratio of the MIC values ($\text{MIC}_{\text{DDS}} = 16 \mu\text{mol/l}$; $\text{MIC}_{\text{SMZ}} = 0.8 \mu\text{mol/l}$; *E. coli*). Also, the synergistic behavior of DDS in combination with a dihydrofolate inhibitor like trimethoprim (TMP) or pyrimethamine (²⁶) is comparable to the




FIG. 12. Example of quantitative relation between apparent *E. coli* growth rate constants k_{app} and DDS concentration (total counts). The curve is plotted in accordance with the equation: $1/(k_0 - k_{\text{app}}) = \frac{1}{C} K_A + K_B$.

synergism observed in the case of SA in combination with TMP (Fig. 13). The mechanism of the synergism of combinations of SA and TMP or DDS and TMP is still not understood. As a hypothesis especially to explain the extraordinarily high antibacterial effect of DDS alone against *M. leprae*, we would propose that the formed DDS-dihydroptericoic acid analog might act itself at the dihydrofolate reductase level, thus causing a more than additive (i.e., synergistic) effect to the inhibitory power of dapsone alone.

Poe (¹⁹) has also made a proposal to explain the synergism observed for combinations of SA and dihydrofolate reductase inhibitors. He assumes that SA and also DDS act not only at the enzyme dihydroptericoic acid synthetase (Fig. 1) but also at the dihydrofolate reductase. The published K_i values (see Table 3) for various SA and DDS acting on dihydrofolate reductase show, however, that they are at least two orders of magnitude larger than the inhibition constants observed for inhibition of




FIG. 13. Typical generation rate curves of *E. coli* (mutafloor) at 37°C in the presence of DDS and TMP alone and in combination. The curves, final drug concentrations ($\mu\text{mol/l}$) and generation rate constants $k_{\text{app}} \times 10^{-4} [\text{sec}^{-1}]$ were as follows: (x) control 3.766; (●) 10 DDS, 3.096; (○) 0.4 TMP, 3.25; (△) 10 DDS, 0.4 TMP \pm 0.0.

dihydropterotic acid synthetase. It is, however, interesting to note that DDS shows a higher affinity towards *E. coli* dihydrofolate reductase than SA whereas the affinity for the main target enzyme dihydropterotic acid synthetase, is opposite as shown above. One could speculate, therefore, that in the case of *M. leprae* the affinity of DDS towards dihydrofolate reductase might even be larger, thus causing an inhibiting effect in addition to the inhibition of dihydropterotic acid synthetase. The two phases of inhibition of DDS seen in *E. coli* cultures could also indicate that there is an inhibitory effect of the analog formed by DDS and phosphorylated dihydropteridine alcohol (see reaction scheme). An experiment was designed in which PABA was added after 4 hr to a culture inhibited by DDS. After this time a certain amount of the DDS analog has been formed; the in-




FIG. 14. Typical generation rate curves of *E. coli* (mutafloor) at 37°C in the presence of DDS, PABA and TMP alone and in combination.

hibitory action of this analog should at least not immediately be antagonized by the addition of PABA, which has no affinity for the dihydrofolate reductase, and could show a synergistic or additive effect in combination with TMP. The results are given in Fig. 14 together with the curves obtained in the presence of DDS alone and in a mixture with TMP. It is clearly demonstrated that at least in the case of *E. coli*, the DDS inhibitory activity can be totally antagonized by PABA, i.e., there seems to be no additional activity of the DDS analog or DDS alone at the dihydrofolate reductase level.

Hypothesis about an additional mechanism of action of DDS in *M. leprae*. It seems reasonable to assume that DDS acts on the folate synthesizing pathway of *M. leprae* in a similar way as shown for *E. coli*, especially if one considers the results obtained for other bacterial strains and even for plasmodia. The reported extremely high inhib-

TABLE 3.

Inhibition constants of sulfonamide for *E. coli* dihydrofolate reductase and *E. coli* dihydropteroate synthetase

Compound	K_1 (M)		K_1^2
	Dihydro- pteroate synthetase	Dihydro- folate reductase	
Sulfaquinoxaline		1.3×10^{-4}	$1.7 \pm 0.4 \times 10^{-4}$
4,4'-Diaminodiphenyl-sulfone		1.8×10^{-4}	$3.0 \pm 1.1 \times 10^{-4}$
Sulfamerazine	4.0×10^{-6}	4.0×10^{-4}	$6.3 \pm 2.2 \times 10^{-4}$
Sulfapyridine	1.3×10^{-5}	5.7×10^{-4}	$1.7 \pm 1.2 \times 10^{-3}$
Sulfathiazole	2.3×10^{-6}	7.4×10^{-4}	$9.6 \pm 4.8 \times 10^{-4}$
Sulfamethoxazole		2.0×10^{-3}	$2.3 \pm 0.3 \times 10^{-3}$
Sulfinilic acid	8.3×10^{-6}	3.5×10^{-3}	$6.5 \pm 0.8 \times 10^{-3}$
Sulfaguanidine	1.7×10^{-5}	3.9×10^{-3}	5.6×10^{-1}
Sulfisoxazole		6.5×10^{-3}	$3.0 \pm 0.3 \times 10^{-3}$
Sulfanilamide	6.7×10^{-5}	2.4×10^{-2}	$1.3 \pm 0.7 \times 10^{-2}$

From: M.Poe: Science 194, 533 (1976)

itory power of DDS against *M. leprae* might be explainable by assuming one or a combination of the following possibilities:

- 1) a very high affinity to the enzyme dihydropteroic acid synthetase. This does not seem to be very likely from the experience with reversible competitive inhibitors. In all cases known so far, the affinity constant of a strong inhibitor was in the order of magnitude of the natural metabolite.
- 2) a very low concentration of the competing PABA within the bacterial cell. Since the MED values of DDS against *M. leprae* are extraordinarily low, this is not very likely.
- 3) an additional inhibitory effect of DDS or the formed analog at the dihydrofolate reductase site or 4) a synergistic effect of DDS or the formed analog at the dihydrofolate reductase. These arguments are still likely to have some relevance even if no evidence can be derived from the studies on *E. coli* as discussed above. Studies on *M. lufu*

with a high sensitivity for DDS are in progress in our laboratory to further address these points.

- 5) accumulation of DDS within bacterial cells of *M. leprae* (deep compartment). In experiments with *M. kansasii*, Panitch and Levy found a 14-15-fold accumulation of DDS within the bacterial cells after 8 days of treatment of the bacterial culture (¹⁷).
- 6) an additional action outside of the folate synthesizing enzyme system.
- 7) The extremely slow generation rate of *M. leprae* and folate metabolism is the rate determining step. Inhibition of folate metabolism by DDS might be very powerful under these circumstances.

SUMMARY

The antibacterial activity of DDS has been studied in whole cell (*E. coli*), cell-free folate synthesizing enzyme extracts and compared to effects obtained for sulfonamides (SA). It is shown that DDS acts as a synthetase inhibitor in the folate syn-

thesizing enzyme system. DDS reacts with the substrate 7,8-dihydro-6-hydroxymethylpterinopyrophosphate to form a 7,8-dihydropteroic acid analog. Bacterial growth kinetic studies were performed to test for possible synergistic activity of the analog in combination with DDS. Possible reasons for the extremely large inhibitory power of DDS against *M. leprae* are discussed.

RESÚMEN

Se estudió la actividad antibacteriana del DDS en células totales (*E. coli*) y en extractos libres de células que contenían las enzimas necesarias para la síntesis del folato. Los resultados se compararon con los obtenidos con sulfonamidas (SA). Se encontró que el DDS actúa como inhibidor en el sistema enzimático que participa en la síntesis del folato. El DDS reacciona con el substrato 7,8-dihidro-6-hidroximetilpterin-pirofosfato para formar un análogo del ácido 7,8-dihidropteroico. También se hicieron estudios sobre la cinética del crecimiento bacteriano para probar la posible actividad sinérgica del análogo en combinación con el DDS. Se discuten algunas posibles razones para explicar el gran poder inhibitorio del DDS sobre el *M. leprae*.

RESUMÉ

L'activité anti-bactérienne de la DDS a été étudiée sur des cellules entières de *E. coli*, ainsi que sur des extraits enzymatiques dépourvus de cellules et synthétisant le folate. Cette activité a été comparée aux effets obtenus avec les sulfonamides. On a ainsi montré que la DDS agit comme inhibiteur de la synthétase, dans le système enzymatique synthétisant le folate. La DDS réagit avec le 7,8-dihydro-6-hydroxymethylpterinopyrophosphate, pour former un analogue de l'acide 7,8-dihydropteroïque. Des études cinétiques de la croissance bactérienne ont été menées afin d'étudier l'activité synergique éventuelle de cet analogue en combinaison avec la DDS. Les raisons qui pourraient expliquer le pouvoir inhibiteur extrêmement puissant de la DDS contre *M. leprae*, sont discutées.

Acknowledgement. This investigation received financial support from the Chemotherapy of Leprosy (THELEP) component of the UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases. It has also been supported by the German Leprosy Relief Association, D-7800 Würzburg, Germany.

REFERENCES

- ANTON, A. H. The relationship between the binding of sulfonamides to albumin and their antibacterial efficacy. *J. Pharmacol. Exp. Ther.* **129** (1960) 282–290.
- BOCK, L., BUTTE, W., RICHTER, M. and SEYDEL, J. K. A simple and rapid thin-layer chromatography method to follow the kinetics of product formation by 7,8-dihydro-6-hydroxymethylpterinpyrophosphokinase and 7,8-dihydropteroate synthetase. *Anal. Biochem.* **86** (1978) 238–251.
- BOCK, L., BUTTE, W., RICHTER, M. and SEYDEL, J. K. Sulfonamide structure-activity relationship in a cell-free system. II. Proof for the formation of a sulfonamide-containing folate analogue. *J. Med. Chem.* **17** (1974) 23–28.
- BROWN, G. M. Methods for measuring inhibition by sulfonamides of the enzymatic synthesis of dihydropteroic acid. *Methods Med. Res.* **10** (1964) 233–238.
- BROWN, G. M. The biosynthesis of folic acid. II. Inhibition by sulfonamides. *J. Biol. Chem.* **237** (1962) 536–540.
- FERONE, R. The enzymatic synthesis of dihydropteroate and dihydrofolate by *Plasmodium berghei*. *J. Protozool.* **20** (1973) 459–464.
- GARRETT, E. R. Classification and evaluation of combined antibiotic activity. *Antibiot. Chemother.* **8** (1958) 8–20.
- IWAI, K. and OKINAKA, O. The biosynthesis of folic acid compounds in plants. II. Some properties of dihydropteroate-synthesizing enzyme in pea seedlings. *J. Vitaminol.* **14** (1968) 170–177.
- JAENICKE, L. and CHAN, P. C. Die Biosynthese der Folsäure. *Angew. Chem.* **72** (1960) 752–753.
- KRÜGER-THIEMER, E., WEMPE, E. and TÖPFER, M. Die antibakterielle Wirkung des nicht eiweißgebunden Anteils der Sulfanilamide in menschlichem Plasmawasser. *Arzneim. Forsch* **15** (1965) 1309–1317.
- LEVY, L. and PETERS, J. H. Some characteristics of the action of dapsone on multiplication of *Mycobacterium leprae* in the mouse. *Lepr. Rev.* **48** (1977) 237–245.
- LEVY, L. and PETERS, J. H. Susceptibility of *Mycobacterium leprae* to dapsone as a determinant of patient response to acedapsone. *Antimicrob. Agents Chemother.* **9** (1976) 102–112.
- MAREN, T. H. Relations between structure and biological activity of sulfonamides. *Annu. Rev. Pharmacol. Toxicol.* (1976) 309–327.
- MILLER, G. H., DOUKAS, P. H. and SEYDEL, J. K. Sulfonamide structure-activity relationships in a cell-free system. Correlation of inhibition of folate synthesis with antibacterial activity and physico-chemical parameters. *J. Med. Chem.* **15** (1972) 700–706.
- MITSUDA, H. and SUZUKI, Y. Enzymatic pyrophosphorylation of 2-amino-4-hydroxy-6-hydroxymethyl-7,8-dihydropteridine by cell-free extracts of *Escherichia coli*. *Biochem. Biophys. Res. Commun.* **36** (1969) 1–6.
- ORTIZ, P. J. and HOTCHKISS, R. D. The enzymatic synthesis of dihydrofolate and dihydropteroate in cell-free preparations from wild-type and sulfonamide-resistant pneumococcus. *Biochemistry* **5** (1966) 67–74.

17. PANITCH, M. L. and LEVY, L. The action of dapsone on a susceptible strain of *Mycobacterium kansasii*. Lepr. Rev. **49** (1978) 131-140.
18. PETERS, J. H., GORDON, G. R., MURRAY, J. F., FIELDSTEEL, A. H., and LEVY, L. Minimal inhibitory concentration of dapsone for *Mycobacterium leprae* in rats. Antimicrob. Agents Chemother. **8** (1975) 551-557.
19. POE, M. Antibacterial synergism: a proposal for chemotherapeutic potentiation between trimethoprim and sulfamethoxazole. Science **194** (1976) 533-535.
20. ROGERS, E. F., CLARK, R. L., BECKER, H. J., PESSOLANO, A. A., LEANZA, W. J., MCMANUS, E. C., ANDRIULI, F. J. and CUCKLER, A. C. Anticoccidial activity of 4-amino-2-ethoxybenzoic acid and related compounds (29616). Proc. Soc. Exp. Biol. Med. **117** (1964) 488-492.
21. SAXENA, A. K. and SEYDEL, J. K. (unpublished results, 1978).
22. SEYDEL, J. K. Prediction of the *in vitro* activity of sulfonamides synthesized from simple amines by use of electronic data obtained from the simple amines. J. Med. Chem. **14** (1971) 724-729.
23. SEYDEL, J. K. Prediction of *in vitro* activity of sulfonamides, using Hammett constants or spectrophotometric data of the basic amines for calculation. Med. Pharmacol. **2** (1966) 259-265.
24. SEYDEL, J. K. The relationship of pharmacokinetic properties of some sulfonamides to certain physicochemical parameters. Int. Congr. Chemother., Proc. I/II (Tokyo) (1970) 881.
25. SEYDEL, J. K., MILLER, G. H. and DOUKAS, P. H. Structure-activity correlations of sulfonamides in cell-free systems compared to correlations obtained in whole cell systems and *in vivo*: In: *Medicinal Chemistry*, P. Pratesi, ed., London: Butterworths, 1973, pp. 139-151.
26. SEYDEL, J. K. and WEMPE, E. Kinetics and mechanism of action of folate synthesis inhibitors, alone or in combination, on *Escherichia coli*. III. Pyrimethamine, trimethoprim and sulfamethoxazole. Chemotherapy **21** (1975) 131-145.
27. SEYDEL, J. K., WEMPE, E., MILLER, G. H. and MILLER, L. Kinetics and mechanism of action of trimethoprim and sulfonamides, alone or in combination upon *E. coli*. Chemotherapy **17** (1972) 217-258.
28. SHEPARD, C. C. Further experience with the kinetic method for the study of drugs against *Mycobacterium leprae* in mice. Activities of DDS, DFD, ethionamide, capreomycin and PAM 1392. Int. J. Lepr. **37** (1969) 389-397.
29. SHIOTA, T., DISRAELY, M. N. and MCCANN, M. P. The enzymatic synthesis of folate-like compounds from hydroxymethyl-dihydropteridine pyrophosphate. J. Biol. Chem. **239** (1964) 2259-2266.
30. WEBB, J. L. *Enzyme and Metabolic Inhibitors, Vol. 1*, New York: Academic Press, Inc., 1963, p. 86.
31. WOLF, B. and HOTCHKISS, R. D. Genetically modified folic acid synthesizing enzymes of pneumococcus. Biochemistry **2** (1963) 145-150.