

- rium leprae* and other mycobacteria. Infect. Immun. **51** (1986) 157–162.
19. SAITO, H., TOMIOKA, H., WATANABE, T. and SATO, K. Mechanism of phagocytosis of *Mycobacterium leprae* and other mycobacteria by human oligodendroglial cells. Infect. Immun. **51** (1986) 163–167.
 20. SPENCER, P. S. and THOMAS, P. K. Ultrastructural studies of the dying-back process. II. The sequestration and removal by Schwann cells and oligodendrocytes of organelles from normal and diseased axons. J. Neurocytol. **3** (1974) 763–783.
 21. THOMAS, P. K. Nerve injury. In: *Essays on the Structure and Function of the Nervous System*. Belairs, R. and Gray, E. G., eds. Oxford: Clarendon Press, 1974, pp. 44–70.
 22. URICH, H. Diseases of peripheral nerves. In: *Greenfield's Neuropathology*. Blackwood, W. and Corsellis, J. A. N., eds. Chicago: Year Bk. Medical Pubs. Inc., 1976, pp. 688–770.
 23. WEISS, P. and WANG, H. Transformation of adult Schwann cells into macrophages. Proc. Soc. Exp. Biol. Med. **58** (1945) 273–275.

Microbe Dependence of *Mycobacterium leprae*: A Possible Intracellular Relationship with Protozoa

TO THE EDITOR:

Kato has postulated that *Mycobacterium leprae* may be dependent on concomitant microbes for its growth^(3,4). We offer an extension of this hypothesis by postulating that the original host of this mycobacterium may have been an amoeba or other free-living protozoan. The human (and armadillo) macrophages, which have certain features in common with amoebae, may thus be alternative host cells. This possibility was raised by one of us (TR) who was intrigued by the paradox that *Legionella pneumophila* has very fastidious growth requirements yet lives in water. Investigations showed that this bacterium survived and multiplied within amoebae⁽⁷⁾. Of particular relevance is the fact that *L. pneumophila* replicates within human macrophages, and parallels have been drawn between infection by this pathogen in immunocompromised patients and lepomatous leprosy⁽¹⁾. Further, the more fastidious *Legionella micdadei* (Pittsburgh pneumonia agent) can in tissue be as acid-fast as *M. leprae*⁽⁵⁾, suggesting a behavioral similarity.

We studied the interaction at 30°C between *M. avium* and an endosymbiont-carrying strain of *Acanthamoeba polyphaga* (Ap-1) which has previously been successfully used to isolate, via amoebal enrichment, *L. pneumophila* serogroup-1, *L. pneumophila* serogroup-3 and a *Legionella*-like amoebal pathogen (LLAP-3) from hu-

man cases of pneumonia⁽⁷⁾. Endosymbionts are bacteria that lie within the amoebal cytoplasm, replicate together with the host cell, and cannot be cultivated *in vitro*, indicating a close mutual dependence⁽⁶⁾. Intriguingly, some human sera contain antibody to the small, curved, gram-negative, non-acid-fast endosymbiont.

A PYG broth culture of the amoebae was inoculated with a recent clinical isolate of *M. avium* in a bacilli-to-amoeba ratio of 10:1. Half of the mycobacteria were phagocytosed within 1 hr, and virtually all within 18 hr. After 3 weeks, all amoebae contained clumps of strongly staining, intact, acid-fast bacilli. The amoebae were subcultured every 3 weeks to a total of seven subcultures, and were found to replicate at the same rate as uninfected controls. During this time, all amoebae contained the clumps of acid-fast bacilli. Amoebae that became distended with bacilli appeared to be able to liberate these into the medium.

These studies showed that *M. avium* was readily phagocytosed by the amoebae, and appeared to establish a relationship with the host cell. This leads us to postulate that *M. leprae* may likewise exist, or have evolved, in close association with protozoa, and we suggest three possibilities. First, *M. leprae* may replicate within certain species or strains of amoebae, as previously suggested by Jadin⁽²⁾. Second, the intracellular growth of *M. leprae* may be dependent on the pres-

ence of endosymbionts. Third, *M. leprae* may itself be an endosymbiont.

In view of these possibilities, it would be of interest to examine protozoa in those environments where the earth-burrowing armadillos have been infected with *M. leprae*, as well as to attempt the infection of various protozoa with this pathogen in the hope of developing a "cell line" for drug-resistance studies and large-scale propagation.

—John M. Grange, M.Sc., M.D.

—C. Ann Dewar, M.I.Biol.

Cardiothoracic Institute

University of London

Fulham Road

London SW3 6HP, U.K.

—Timothy J. Rowbotham, B.Sc., Ph.D.

Public Health Laboratory

Bridle Path

York Road

Leeds LS15 7TR, U.K.

REFERENCES

1. BHARDWAJ, N., NASH, T. W. and HORWITZ, M. A. Interferon- γ -activated human monocytes inhibit the intracellular multiplication of *Legionella pneumophila*. *J. Immunol.* **137** (1986) 2662–2669.
2. JADIN, J. B. Amibies "limax" vecteurs possibles des mycobacteries et de *Mycobacterium leprae*. *Acta Leprol.* **59-60** (1975) 57–66.
3. KATO, L. Absence of mycobactin in *Mycobacterium leprae*; probably a microbe dependent microorganism. Implications. *Indian J. Lepr.* **57** (1985) 58–70.
4. KATO, L. A multifactorial culture medium with growth factors from leprosy-derived mycobacteria proposed in cultivation trials for *Mycobacterium leprae*. *Int. J. Lepr.* **54** (1986) 310–311.
5. MYEROWITZ, R. L., PASCULLE, A. W., DOWLING, J. N., PAZIN, G. J., PUERZER, M., YEE, R. E., RINALDO, C. R. and HAKALA, T. R. Opportunistic lung infection due to "Pittsburgh pneumonia agent." *N. Engl. J. Med.* **301** (1979) 953–958.
6. PROCA-CIOBANU, M., LUPASCU, G., PETROVICI, A. and IONESCU, M. D. Electron microscopic study of a pathogenic *Acanthamoeba castellanii* strain: the presence of bacterial endosymbionts. *Int. J. Parasitol.* **5** (1975) 49–56.
7. ROWBOTHAM, T. J. Isolation of *Legionella pneumophila* from clinical specimens via amoebae, and the interaction of those and other isolates with amoebae. *J. Clin. Pathol.* **36** (1983) 978–986.
1. BHARDWAJ, N., NASH, T. W. and HORWITZ, M. A. Interferon- γ -activated human monocytes inhibit the

Deficiency in the Biosynthesis of Interleukin 2 (IL-2) and Functional Presence of the IL-2 Receptor in Lepromatous Leprosy

TO THE EDITOR:

Patients with lepromatous leprosy (LL) have a depressed cellular immune capacity to respond to antigens from *Mycobacterium leprae*, a reduction in effector and helper T cells, and an increase in suppressor T cells (2, 4, 11, 16). A reduction in the synthesis of interleukins has been postulated in some murine models (9). In man, it has been shown that upon the addition of interleukin 2 (IL-2) to cultures of lymphocytes from patients with LL, these lymphocytes recover their capacity to proliferate (7, 8). However, in another paper (12) it has been pointed out that this phenomenon does not occur.

We report here the results of a study looking for the production of IL-2 by lymphocytes from patients with LL, and the func-

tional presence of IL-2 receptors on their lymphocytes.

Study subjects. Twenty patients, 11 men and 9 women, from the Instituto Dermatológico at Guadalajara, Jalisco, México, were diagnosed as having nodular LL according to international criteria (14). All presented positive bacilloscopy, and all received an irregular treatment of 100 mg of dapsone (diaminodiphenyl sulfone) per day. The length of treatment ranged from 2 to 15 years. Fourteen patients were studied for the biosynthesis of IL-2 by T lymphocytes obtained from peripheral blood, and the functional presence of the IL-2 receptor on T lymphocytes was studied in 6 patients. Controls consisted of 14 unrelated healthy subjects for the first study and 6 apparently