

detected by the biothesiometer might only be a relatively late change. However, differences in threshold were also noted on those patients who were either untreated or who had been treated for 2 months or less. It is suggested that to investigate this further, loss of vibration sense should be assessed in parallel with other nerve functions (preferably in a quantitative manner) in patients when they first present.

Studies have shown that up to 77% of clinically unaffected nerves in leprosy may show marked impairment of nerve conduction (7). Thus it is of obvious importance to attempt to improve clinical testing in order to assess more subtle levels of nerve damage. This is not only of value in diagnosis but also in follow-up, assessment of treatment and reactions.

We have shown elsewhere that the biothesiometer may be of value in the assessment of insensitive feet and their susceptibility to ulceration and tarsal disintegration (Hammond, C. and Klenerman, P., unpublished data and Klenerman, P., Hammond, C. J., and Kulkarni, V., submitted for publication). It is suggested on the basis of this study that it might also be of use in the diagnosis of the anesthetic patch.

—Paul Klenerman, B.A.

—Christopher Hammond, B.A.

*Oxford University Medical School  
Oxford, England*

**Acknowledgments.** We would like to thank Dr. Jal Mehta (Hon. Pres.) and all of the staff at the Dr. Bendorawalla Leprosy Hospital, Pune; Dr. Sunil Tolat, Dermatology Department, Sassoon Hospital, Pune; and Dr. T. J. Ryan, The Slade Hospital, Oxford, for their help and support with this project. We would also like to thank Dr. Colin McDougall and Prof. L. Klenerman for providing us with ideas, equipment, and encouragement.

We are grateful for grants from LEPRO, the *Journal of Bone and Joint Surgery*, St. Francis Leprosy Guild, Medical Defence Union, Anglo-American (Zimbabwe) and the Oxford University Medical School.

#### REFERENCES

1. CHATTOPADHYAY, S. P., BONSA, P. C. and RATHORE, B. S. Value of pilocarpine test in early diagnosis of leprosy. *Indian J. Lepr.* **56** (1984) 877–883.
2. FOX, J. C. and KLEMPERER, W. Vibratory sensibility; a quantitative study of its thresholds in nervous disorders. *Arch. Neurol. Psychiat.* **48** (1942) 622–645.
3. GRAY, R. C. Quantitative study of vibration sense in normals and pernicious anemia cases. *Minnesota Med.* **15** (1932) 674. (Quoted in Ref. 4).
4. GREGG, E. C. Absolute measurement of vibratory threshold. *Arch. Neurol. Psychiat.* **66** (1951) 403–411.
5. IGGO, A. Cutaneous sensory mechanisms. In: *The Senses*. Barlow, H. D. and Mollon, J. D., eds. Cambridge: Cambridge University Press, 1982, pp. 389–408.
6. JAIN, G. L., PASRICHA, J. S. and GUHA, S. K. Minimum temperature felt as hot (MTH)—a new concept for grading the loss of temperature sensation in leprosy patients. *Int. J. Lepr.* **53** (1985) 206–210.
7. McLeod, J. G., Hargrave, J. C., Walsh, J. C., Booth, G. C., Gye, R. S. and BARRON, A. Nerve conduction studies in leprosy. *Int. J. Lepr.* **43** (1975) 21–31.
8. PFALTZGRAFF, R. E. and BRYCESON, A. Clinical leprosy. In: *Leprosy*. Hastings, R. C., ed. Edinburgh: Churchill Livingstone, 1985, pp. 134–176.
9. WEDDELL, A. G. M., JAMISON, D. G. and PALMER, E. Recent investigations into the sensory and neurohistological changes in leprosy. In: *Leprosy in Theory and Practice*. Cochrane, R. G. and Davey, T. F., eds. Bristol: John Wright & Sons Ltd., 1964, pp. 205–220.
10. WILLIAMS, G., GILL, J. S., ABER, V. and MATHER, H. M. Variability in vibration perception threshold among sites; a potential source of error in biothesiometry. *Br. Med. J.* **296** (1988) 233–235.

## Isolation of Mycobacteria from Healthy Persons' Stools

TO THE EDITOR:

In 1986, Collins published in this JOURNAL a paper “*Mycobacterium avium*-complex infections and development of the acquired immunodeficiency syndrome: ca-

sual opportunist or causal cofactor?” (4) One question brought up in his paper concerned the possible gastrointestinal origin of *Mycobacterium avium-intracellulare* (MAI) infections in some acquired immunodeficien-

cy syndrome (AIDS) patients. Indeed, the MAI complex is frequently a cause of disseminated infection in patients with AIDS (19), and the isolation of MAI from stools of AIDS patients has been reported (1, 6, 8, 10, 14, 21-23, 28). However, little is known about the presence of mycobacteria in the stools of healthy individuals. We therefore decided to study the mycobacterial flora of the stools of 50 healthy European volunteers and to identify the mycobacteria by both conventional tests and by lipid analysis. All specimens were examined by microscopy after Ziehl-Neelsen staining. The mycobacteria were isolated after decontaminating the specimens as described by Portaels, *et al.* (20), i.e., a modified procedure of Wolinsky and Rynearson (27). The decontaminated samples were inoculated onto four tubes of Ogawa egg yolk medium, incubated at 30°C for 6 months, and observed for growth every week. When a tube was suspected of being positive for mycobacteria, each colony was subcultured on Ogawa medium and checked for acid fastness by Ziehl-Neelsen staining. The mycobacteria were identified according to well-established methods (12, 25), and their fatty acid and mycolic acid compositions were analyzed by gas chromatography (GC) (13, 15, 16) and thin-layer chromatography (TLC) (5, 7, 18).

No acid-fast bacilli were observed in any of the 50 stool specimens after direct Ziehl-Neelsen staining. Twenty-six specimens (52%) were culture-positive for mycobacteria; the remaining 24 (48%) were negative. The number of colonies observed on the media varied between one and three. Six positive primary cultures failed to grow in subcultures, and from the 20 remaining positive specimens, 26 different mycobacterial strains were cultivated. From 14 specimens only one mycobacterial species was isolated, while from 6 specimens two different species were cultivated (The Table).

Fourteen of the 26 strains were identified as *M. simiae*. They produced  $\alpha$ -,  $\alpha'$ - and ketomycolates like *M. simiae* but were nonchromogenic. Six of the strains were also analyzed by GC; all contained tuberculostearic acid but were devoid of secondary alcohols (2-octadecanol and 2-eicosanol).

Five strains identified as *M. gordonae*

THE TABLE. *Mycobacterial cultures from healthy persons' stools.*

Mycobacterial strains (26)	No. positive specimens (20)
<i>M. simiae</i>	8
<i>M. gordonae</i>	3
MAI <sup>a</sup>	2
<i>M. malmoense</i>	1
MAI <sup>a</sup> + <i>M. simiae</i>	3
<i>M. gordonae</i> + <i>M. simiae</i>	2
<i>M. malmoense</i> + <i>M. simiae</i>	

<sup>a</sup> MAI = *M. avium-intracellulare* complex.

contained  $\alpha$ -, methoxy- and ketomycolates. The presence of a branched chain C14 and a hydroxylated C20 acid as well as the absence of tuberculostearic acid and secondary alcohols—all characteristic features of this species—were established by GC analysis.

Five nonchromogenic strains were classified as belonging to the MAI complex. The results of the TLC studies revealed the presence of  $\alpha$ -, keto- and  $\omega$ -carboxymycolates. Tuberculostearic acid and the secondary alcohols were identified by GC.

Two nonchromogenic strains which both grew poorly at 37°C were identified as *M. malmoense* according to the results of the TLC and GC studies. Their mycolate patterns were identical to those of *M. malmoense* and *M. simiae* ( $\alpha$ -,  $\alpha'$ - and ketomycolates), but, uniquely for *M. malmoense*, both strains contained appreciable amounts of 2-methyl eicosanoic and 2,4,6-trimethyl tetracosanoic acid (24).

The isolation of mycobacteria from human intestinal tissues (2, 3, 9) and from AIDS patients' stools (1, 6, 8, 10, 14, 21-23, 28) has been reported repeatedly. While a variety of different species (MAI, *M. fortuitum* complex, *M. chelonae*) have been found in intestinal specimens from patients with Crohn's disease, ulcerative colitis, and noninflammatory bowel diseases (2, 3, 9), the predominance of MAI strains from intestinal tissues and stools of AIDS patients is striking (1, 6, 8, 10, 14, 21-23, 28). The present study shows that mycobacteria can also be isolated from more than 50% of the stools from healthy individuals. We found a predominance of *M. simiae* followed by MAI strains, *M. gor-*

*donae*, and *M. malmoense*. Our findings of *M. simiae* and *M. malmoense*, both recognized as causes of human diseases (<sup>11, 17, 26</sup>), are particularly noteworthy. Environmental isolates of these species have not been reported although the epidemiology of the diseases for which they were responsible strongly suggests their presence in the environment. The mycobacteria might enter the host via contaminated food or water and then colonize the gastrointestinal tract. As suggested by Collins (<sup>4</sup>), it is possible that some of these mycobacteria (e.g., MAI) possess factors which enable them to attach to the intestinal mucosae, colonize the membranes, and invade them when the AIDS virus has depleted the T-cell defenses. To evaluate whether the mycobacteria isolated from stools originate from an occasional colonization of the intestinal tract or whether they form a part of the permanent flora of some individuals, samples should regularly be studied from the same subjects on a long-term basis. Additional studies on stools from healthy persons, from persons at high risk for developing AIDS, and from AIDS patients with and without disseminated MAI infections are also required for confirmation of the hypothesis that intestinal colonization precedes mycobacterial infection.

—Françoise Portaels, Ph.D.

Associate Professor  
Department of Microbiology  
Chief, Laboratory of Mycobacteriology  
Institute of Tropical Medicine  
Nationalestraat 155  
B-2000 Antwerp, Belgium

—Lennart Larsson, Ph.D.

Associate Professor  
Department of Medical  
Microbiology  
University of Lund  
Sölvegatan 23  
S-233 62 Lund, Sweden

—Pascale Smeets

Laboratory Technician  
Laboratory of Mycobacteriology  
Department of Microbiology  
Institute of Tropical Medicine  
Nationalestraat 155  
B-2000 Antwerp, Belgium

## REFERENCES

1. ARMSTRONG, D. A., GOLD, J. W., DRYJANSKI, J., WHIMBEY, E., POLSKY, B., HAWKINS, C., BROWN, A. E., BERNARD, E. and KIEHN, T. E. Treatment of infections in patients with the acquired immunodeficiency syndrome. *Ann. Intern. Med.* **103** (1985) 738–743.
2. CHIODINI, R. J., VAN KRUIJNINGEN, H. J., MERKAL, R. S., THAYER, W. R., JR. and COUTU, J. A. Characteristics of an unclassified *Mycobacterium* species isolated from patients with Crohn's disease. *J. Clin. Microbiol.* **20** (1984) 966–971.
3. COLEMONT, L. J., PATTYN, S. R., MICHIENSEN, P. P., PEN, J. H., PELCKMANS, P. A., VAN MAERCKE, Y. M. and PORTAELS, F. Acid-fast bacilli in Crohn's disease. *Lancet* **1** (1988) 294–295.
4. COLLINS, F. M. *Mycobacterium avium*-complex infections and development of the acquired immunodeficiency syndrome: casual opportunist or causal cofactor? *Int. J. Lepr.* **54** (1986) 458–474.
5. DAFFÉ, M., LANÉLLE, M. A., ASSELINEAU, C., LÉVY-FRÉBAULT, V. and DAVID, H. L. Intérêt taxonomique des acides gras des mycobactéries: proposition d'une méthode d'analyse. *Ann. Microbiol.* **134B** (1983) 241–256.
6. DAMSKER, B. and BOTTONNE, E. J. *Mycobacterium avium-Mycobacterium intracellulare* from the intestinal tracts of patients with the acquired immunodeficiency syndrome: concepts regarding acquisition and pathogenesis. *J. Infect. Dis.* **151** (1985) 179–181.
7. DOBSON, G., MINNIKIN, D. E., MINNIKIN, S. M., PARLETT, J. M., GOODFELLOW, M., RIDELL, M. and MAGNUSSON, M. Systematic analysis of complex mycobacterial lipids. In: *Chemical Methods in Bacterial Systematics*. Goodfellow, M. and Minnikin, D. E., eds. London: Academic Press, 1985, pp. 237–265.
8. GILLIN, J. S., URMACHER, C., WEST, R. and SHIKE, M. Disseminated *Mycobacterium avium-intracellulare* infection in acquired immunodeficiency syndrome mimicking Whipple's disease. *Gastroenterology* **85** (1983) 1187–1191.
9. GRAHAM, D. Y., MARKESICH, D. C. and YASHINURA, H. Mycobacteria and inflammatory bowel disease. *Gastroenterology* **92** (1987) 436–442.
10. HAWKINS, C. C., GOLD, J. W. M., WHIMBEY, E., KIEHN, T. E., BRANNON, P., CAMMARATA, R., BROWN, A. E. and ARMSTRONG, D. *Mycobacterium avium* complex infections in patients with the acquired immunodeficiency syndrome. *Ann. Intern. Med.* **105** (1986) 184–188.
11. JENKINS, P. A. *Mycobacterium malmoense*. *Tubercle* **66** (1985) 193–195.
12. JENKINS, P. A., PATTYN, S. R. and PORTAELS, F. Diagnostic bacteriology. In: *The Biology of the Mycobacteria*. Ratledge, C. and Stanford, J. L., eds. London: Academic Press, 1982, pp. 441–471.

13. JIMENEZ, J. and LARSSON, L. Heating cells in acid methanol for 30 min without freeze-drying provides adequate yields of fatty acids and alcohols for gas chromatographic characterization of mycobacteria. *J. Clin. Microbiol.* **24** (1986) 844–845.
14. KIEHN, T. E., EDWARDS, F. F., BRANNON, P., TSANG, A. Y., MAIO, M., GOLD, J. W. M., WHIMBEY, E., WONG, B., McCLATCHY, J. K. and ARMSTRONG, D. Infections caused by *Mycobacterium avium* complex in immunocompromised patients: diagnosis by blood culture and fecal examination, antimicrobial susceptibility tests, and morphological and seroagglutination characteristics. *J. Clin. Microbiol.* **21** (1985) 168–173.
15. LARSSON, L., DRAPER, P. and PORTAELS, F. Use of gas chromatography to differentiate *Mycobacterium leprae* from cultivable armadillo-derived mycobacteria, *M. avium/intracellulare*, and *M. lepraemurium* by analysis of secondary alcohols. *Int. J. Lepr.* **53** (1985) 441–446.
16. LARSSON, L. and ODHAM, G. Injection principles on capillary gas chromatographic analysis of bacterial fatty acids. *J. Microbiol. Meth.* **3** (1984) 77–82.
17. LÉVY-FRÉBAULT, V., PANGNON, B., BURÉ, A., KATLAMA, C., MARCHE, C. and DAVID, H. L. *Mycobacterium simiae* and *Mycobacterium avium-M. intracellulare* mixed infection in acquired immune deficiency syndrome. *J. Clin. Microbiol.* **25** (1987) 154–157.
18. MINNIKIN, D. E., MINNIKIN, S. M., PARLETT, J. M., GOODFELLOW, M. and MAGNUSSON, M. Mycolic acid patterns of some species of *Mycobacterium*. *Arch. Microbiol.* **139** (1984) 225–231.
19. PORTAELS, F. Le SIDA et les mycobactéries atypiques. *Ann. Soc. Belg. Med. Trop.* **67** (1987) 93–116.
20. PORTAELS, F., DE MUYNCK, A. and SYLLA, M. P. Selective isolation of mycobacteria from soil: a statistical analysis approach. *J. Gen. Microbiol.* **134** (1988) 849–855.
21. ROTH, R. I., OWEN, R. L., KEREN, D. F. and VOLBERDING, P. A. Intestinal infection with *Mycobacterium avium* in acquired immune deficiency syndrome (AIDS). *Dig. Dis. Sci.* **30** (1985) 497–504.
22. SCHNEEBAUM, C. W., NOVICK, D. M., CHABON, A. B., STRUTYNSKY, N., YANCOVITZ, R. and FREUND, S. Terminal ileitis associated with *Mycobacterium avium-intracellulare* infection in a homosexual man with acquired immune deficiency syndrome. *Gastroenterology* **93** (1987) 1127–1132.
23. STACEY, A. R. Isolation of *Mycobacterium avium-intracellulare-scrofulaceum* complex from faeces of patients with AIDS. *Br. Med. J.* **293** (1986) 1194.
24. VALERO-GUILLEN, P., MARTIN-LUENGO, F., LARSSON, L., JIMENEZ, J., JUHLIN, I. and PORTAELS, F. Fatty and mycolic acids of *Mycobacterium malmoense*. *J. Clin. Microbiol.* **26** (1988) 153–154.
25. WAYNE, L. G. The atypical mycobacteria: recognition and disease association. *Crit. Rev. Microbiol.* **12** (1985) 185–222.
26. WOLINSKY, E. Nontuberculous mycobacteria and associated diseases. *Am. Rev. Respir. Dis.* **119** (1979) 107–159.
27. WOLINSKY, E. and RYNEARSON, T. K. Mycobacteria in soil and their relation to disease-associated strains. *Am. Rev. Respir. Dis.* **97** (1968) 1032–1037.
28. YOUNG, L. S., INDERLIED, C. B., BERLIN, O. B. and GOTTLIEB, M. S. Mycobacterial infections in AIDS patients with an emphasis on the *Mycobacterium avium* complex. *Rev. Infect. Dis.* **8** (1986) 1024–1033.

## Comedones Induced by Coconut Oil in a Borderline Tuberculoid Lesion

### TO THE EDITOR:

A 42-year-old female presented with a hypopigmented anesthetic patch over her left knee of 1½ years duration. Diagnosed as a case of tuberculoid leprosy, she had been treated with monthly rifampin and daily dapsone for 13 months. She had been advised to discontinue treatment, and she came to us for a second opinion.

On examination, comedones were observed over the hypopigmented, anesthetic, borderline tuberculoid lesion (Fig. 1). The comedones could be extracted with a comedo extractor. A biopsy stained with hematoxylin and eosin (H&E) showed a compact, perineural and periadnexal epithelioid granuloma in the reticular and deeper der-