

REFERENCES

1. BHATKI, W. S., CHULAWALA, R. G., BAPAT, C. V. and DEO, M. G. Reversal reaction in lepromatous patients induced by a vaccine containing killed ICRC bacilli—a report of five cases. *Int. J. Lepr.* **51** (1984) 466–472.
2. DEO, M. G., BAPAT, C. V., BHALERAO, V., CHATURVEDI, R. M., CHULAWALA, R. G. and BHATKI, W. S. Antileprosy potentials of the ICRC vaccine: a study in patients and healthy volunteers. *Int. J. Lepr.* **51** (1984) 540–549.
3. GROSSINSKY, C. M., JACOB, W. R., CLARK-CURTISS, J. E. and BLOOM, B. R. Genetic relationship among *Mycobacterium leprae*, *Mycobacterium tuberculosis*, and candidate leprosy vaccine strains determined by DNA hybridization: identification of an *M. leprae*-specific repetitive sequence. *Infect. Immun.* **57** (1989) 1535–1541.
4. MUKHERJEE, A., ZAHEER, S. A., SHARMA, A. K., MISRA, R. S., KAR, H. K., MUKHERJEE, R. and TALWAR, G. P. Histopathological monitoring of an immunotherapeutic trial with *Mycobacterium w*. *Int. J. Lepr.* **60** (1992) 28–35.
5. MUSTAFA, A. S. and TALWAR, G. P. Five cultivable mycobacterial strains giving blast transformation and leucocyte migration inhibition of leucocytes analogous to *Mycobacterium leprae*. *Lepr. India* **50** (1978) 498–508.
6. MUSTAFA, A. S. and TALWAR, G. P. Delayed hypersensitivity skin reactions to homologous and heterologous antigens in guinea-pigs immunized with *M. leprae* and four selected cultivable mycobacterial strains. *Lepr. India* **50** (1978) 509–519.
7. SAHIB, H. S. M. and VELLUT, C. Some observations on skin reactions induced by lepromin and four other mycobacterial antigens. *Lepr. India* **50** (1978) 579–587.
8. STANFORD, J. L. Immunotherapy for mycobacterial disease. In: *The Biology of the Mycobacteria. Vol. 3. Clinical Aspects of Mycobacterial Disease*. Ratledge, C., Stanford, J. and Grange, J. M. London: Academic Press, 1989, 567–596.

Dr. Mukherjee, *et al.* Respond to Dr. Deo

TO THE EDITOR:

In response to Dr. Deo's letter claiming that ICRC and *Mycobacterium w* "may not be very different," the following points are offered for consideration.

1) ICRC was isolated by Bapat, *et al.* (1) from leproma nodules and grown in a "conditioned" medium. This organism belongs to the *M. avium-intracellulare* complex and grows in T/900R mice with evidence of dissemination in the liver and sciatic nerve with production of foot drop (2).

2) *Mycobacterium w* has its origin in a collection of atypical mycobacteria grown from sputum specimens and made available in 1974 to Dr. G. P. Talwar by Dr. S. P. Tripathy who was Director of the Tuberculosis Research Center in Madras at that time. This organism, probably a commensal of the upper respiratory tract, is a fast grower and is nonpathogenic to mice, rats and guinea pigs. Its growth and metabolic properties (4,6) are similar to organisms belonging to Runyon's Group 4.

3) The results of DNA hybridization studies on the two organisms reported by Grossinsky, *et al.* (3) show that at a low stringency the percentage of binding of *M. leprae* DNA is 10.5 for ICRC and *Mycobacterium w*, 10.8 for *M. avium* and 10.3 for *M. phlei*. More importantly, at a high stringency *Mycobacterium w* has a distinctly higher binding, i.e., 3.9% as compared to 2.9% for ICRC. In the same article the authors have made the statement that *Mycobacterium w* shows the highest degree of homology with *M. leprae* among the three candidate vaccines.

4) With reference to the RFLP patterns studied by the same workers, the restriction enzymes selected and the regions probed are rather conserved. Restriction and probing of other regions are likely to bring out dissimilarities between the two organisms.

The data from the source, growth characteristics, animal pathogenicity and DNA analysis stated above make it clear that the two organisms are quite different. The therapeutic efficacy obtained with *Mycobacterium w* used in conjunction with multidrug therapy (MDT) has been clearly and unequivocally demonstrated in a number of studies (5,7,8). The only logical way by which the comparative merits of these two different organisms for immunoprophylaxis or immunotherapy can be gauged is with a comparative trial on a standard protocol by an independent party.

We look forward to the results from the ongoing comparative vaccine immunoprophylaxis trial in progress in Avadi, Tamil Nadu, India, where *Mycobacterium w*, ICRC (contributed by Dr. C. V. Bapat), and killed *M. leprae* plus BCG are under a coded comparative trial.

We hope that the data presented above clarifies any doubt about the identity of *Mycobacterium w*.

—Ashok Mukherjee, M.D.
 Syed A. Zaheer, M.B.B.S., D.V.D., Ph.D.
 Amar K. Sharma, M.D.
 Radhey S. Misra, M.D., D.D.
 H. K. Kar, M. D.
 Rama Mukherjee, Ph.D.
 Gurusaran P. Talwar, D.Sc., F.I.C.A.,
 F.A.Sc., F.N.A.

Institute of Pathology-ICMR
Departments of Dermatology
Safdarjang and R.M.L. Hospitals
and the National Institute of
Immunology
New Delhi, India

REFERENCES

1. BAPAT, C. V. and MODAK, M. S. Growth of ICRC bacilli in footpad of mice. *Lepr. India* **50** (1978) 144–155.
2. BAPAT, C. V., RANADIVE, K. J. and KHANOLKAR, V. R. Growth characteristics of an acid-fast mycobacterium isolated from human lepromatous leprosy. *Int. J. Lepr.* **29** (1961) 329–342.
3. GROSSINSKY, C. M., JACOB, W. R., CLARK-CURTISS, J. E. and BLOOM, B. R. Genetic relationship among *Mycobacterium leprae*, *Mycobacterium tuberculosis* and candidate leprosy vaccine strains determined by DNA hybridization: identification of an *M. leprae*-specific repetitive sequence. *Infect. Immun.* **57** (1989) 1535–1541.
4. KATOCH, V. M. A report on the biochemical analysis of *Mycobacterium w*. *Lepr. India* **53** (1981) 385–389.
5. MUKHERJEE, A., ZAHEER, S. A., SHARMA, A. K., MISRA, R. S., KAR, H. K., MUKHERJEE, R. and TALWAR, G. P. Histopathological monitoring of an immunotherapeutic trial with *Mycobacterium w*. *Int. J. Lepr.* **60** (1992) 28–35.
6. SAXENA, V. K., SINGH, U. S. and SINGH, A. K. Bacteriological study of a rapidly growing strain of *Mycobacterium*. *Lepr. India* **50** (1978) 588–596.
7. TALWAR, G. P., ZAHEER, S. A., MUKHERJEE, R., WALIA, R., MISRA, R. S., SHARMA, A. K., KAR, H. K., MUKHERJEE, A., PARIDA, S. K., SURESH, N. R., NAIR, S. K. and PANDEY, R. M. Immunotherapeutic effects of an anti-leprosy vaccine based on a saprophytic cultivable mycobacterium, *Mycobacterium w*, in multibacillary leprosy patients. *Vaccine* **8** (1990) 121–129.
8. TALWAR, G. P., ZAHEER, S. A., SURESH, N. R., PARIDA, S. K., MUKHERJEE, R., SINGH, I. G., SHARMA, A. K., KAR, H. K., MISRA, R. S. and MUKHERJEE, A. Immunotherapeutic trials with a candidate anti-leprosy vaccine based on *Mycobacterium w*. *Trop. Med. Parasitol.* **41** (1990) 369–370.
9. ZAHEER, S. A., MUKHERJEE, R., BEENA, R., MISRA, R. S., SHARMA, A. K., KAR, H. K., KAUR, M., NAIR, S., MUKHERJEE, A. and TALWAR, G. P. Combined multidrug and *Mycobacterium w* in vaccine therapy in patients with multibacillary leprosy. *J. Infect. Dis.* **167** (1993) 401–410.

Leprosy and HIV Infection

TO THE EDITOR:

Leprosy and HIV infection are diseases of an endemic nature in Brazil and are a source of great concern for public health agencies. Brazil is second in the world, after India, in absolute numbers of leprosy cases. According to the Brazilian Health Ministry, the overall prevalence is 4–4.5 cases per 1000 inhabitants and 2 per 1000 in the Ribeirão Preto region. With respect to AIDS, Brazil is among the five countries in the world with the largest absolute number of recorded cases. According to the World Health Organization, the incidence of AIDS

recorded in 1990 was 22.9 per 100,000 inhabitants. According to the Health Department of the state of São Paulo, the incidence of AIDS in the city of Ribeirão Preto is 100.7 per 100,000 inhabitants, and among injectable drug users the HIV prevalence is still 46.5 per 100,000 inhabitants.

The association of HIV infection with infection by mycobacteria such as *Mycobacterium avium* and *M. intracellulare* has been reported (¹). Over the last few years, there have been reports of HIV infection associated with *M. leprae* infection. In the present report, we describe the first case of such